

1) Suppose that

$$\begin{pmatrix} X_1 \\ X_2 \\ X_3 \\ X_4 \end{pmatrix} \sim N_4 \left( \begin{pmatrix} 49 \\ 25 \\ 9 \\ 4 \end{pmatrix}, \begin{pmatrix} 2 & -1 & 3 & 0 \\ -1 & 5 & -3 & 0 \\ 3 & -3 & 5 & 0 \\ 0 & 0 & 0 & 4 \end{pmatrix} \right).$$

a) Find the distribution of  $(X_1, X_3)^T$ .

3.21

$$N_2 \left[ \begin{pmatrix} 49 \\ 9 \end{pmatrix}, \begin{pmatrix} 2 & 3 \\ 3 & 5 \end{pmatrix} \right]$$

b) Which pairs of random variables  $X_i$  and  $X_j$  are independent?

$$X_1 \perp X_4, \quad X_2 \perp X_4, \quad X_3 \perp X_4$$

c) Find the correlation  $\rho(X_1, X_3)$ .

51

$$\frac{\text{Cov}(X_1, X_3)}{\sqrt{\text{Var}(X_1)\text{Var}(X_3)}} = \frac{3}{\sqrt{2 \cdot 5}} = \frac{3}{\sqrt{10}} = 0.9487$$

2) Recall that if  $\mathbf{X} \sim N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ , then the conditional distribution of  $\mathbf{X}_1$  given that  $\mathbf{X}_2 = \mathbf{x}_2$  is multivariate normal with mean  $\boldsymbol{\mu}_1 + \boldsymbol{\Sigma}_{12}\boldsymbol{\Sigma}_{22}^{-1}(\mathbf{x}_2 - \boldsymbol{\mu}_2)$  and covariance matrix  $\boldsymbol{\Sigma}_{11} - \boldsymbol{\Sigma}_{12}\boldsymbol{\Sigma}_{22}^{-1}\boldsymbol{\Sigma}_{21}$ . Let  $Y$  and  $X$  follow a bivariate normal distribution

3.22

$$\begin{pmatrix} Y \\ X \end{pmatrix} \sim N_2 \left( \begin{pmatrix} 49 \\ 17 \end{pmatrix}, \begin{pmatrix} 3 & -1 \\ -1 & 4 \end{pmatrix} \right). \quad \mu_y + \boldsymbol{\Sigma}_{yx}\boldsymbol{\Sigma}_{xx}^{-1}(x - \mu_x)$$

a) Find  $E(Y|X)$ .  $49 + (-1) \frac{1}{4} (x_2 - 17)$  2 Fine

$$= 49 - \frac{1}{4} (x_2 - 17) = 49 + \frac{17}{4} - \frac{1}{4} x_2 = \boxed{53.25 - 0.25x_2}$$

b) Find  $\text{Var}(Y|X)$ .

34

$$3 - (-1) \frac{1}{4} (-1) = 3 - \frac{1}{4} = \boxed{2.75} = \frac{11}{4}$$

→ 3) Suppose  $Y \sim N_n(\mathbf{X}\beta, \sigma^2\mathbf{I})$ . Find the distribution of  $\underline{z} = (\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{Y}$  if  $\mathbf{X}$  is an  $n \times p$  full rank constant matrix and  $\beta$  is a  $p \times 1$  constant vector. *Simplify*

$$E\underline{z} = (\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{X}\beta = \underline{\beta}$$

$$\text{COV}(\underline{z}) = (\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\sigma^2\mathbf{I}\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}$$

3.23

$$= \sigma^2 (\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}$$

$$= \sigma^2 (\mathbf{X}^T\mathbf{X})^{-1}$$

$$\text{So } \underline{z} \sim N_p \left[ \underline{\beta}, \sigma^2 (\mathbf{X}^T\mathbf{X})^{-1} \right]$$

15