

# Multivariate Matching Polynomials of Cyclically Labelled Graphs

John P. McSorley and Philip Feinsilver,  
Department of Mathematics,  
Mailcode 4408,  
Southern Illinois University,  
Carbondale. IL 62901-4408.  
mcsorley60@hotmail.com  
pfeinsil@math.siu.edu

## Keywords

Matching, Matching Polynomial, Labelled, Graph, Recurrence,  
MacMahon's Master Theorem

## Abstract

We consider the matching polynomials of graphs whose edges have been cyclically labelled with the ordered set of  $t$  labels  $\{x_1, \dots, x_t\}$ .

We first work with the cyclically labelled path, with first edge label  $x_i$ , followed by  $N$  full cycles of labels  $\{x_1, \dots, x_t\}$ , and last edge label  $x_j$ . Let  $\Phi_{i,Nt+j}$  denote the matching polynomial of this path. It satisfies the  $(\tau, \Delta)$ -recurrence:  $\Phi_{i,Nt+j} = \tau \Phi_{i,(N-1)t+j} - \Delta \Phi_{i,(N-2)t+j}$ , where  $\tau$  is the sum of all non-consecutive cyclic monomials in the variables  $\{x_1, \dots, x_t\}$  and  $\Delta = (-1)^t x_1 \cdots x_t$ . A combinatorial/algebraic proof and a matrix proof of this fact are given. Let  $G_N$  denote the first fundamental solution to the  $(\tau, \Delta)$ -recurrence. We express  $G_N$  (i) as a cyclic binomial using the Symmetric Representation of a matrix, (ii) in terms of Chebyshev polynomials of the second kind in the variables  $\tau$  and  $\Delta$ , and (iii) as a quotient of two matching polynomials. We extend our results from paths to cycles and rooted trees.

# Introduction

The matching polynomial of a graph is defined in Farrell [1]. Often in pure mathematics and combinatorics it is interesting to consider cyclic structures, *eg.*, cyclic groups, cyclic designs, and circulant graphs. Here we consider the (multivariate) matching polynomial of a graph whose edges have been cyclically labelled.

We concentrate mainly on paths, cycles and trees. To cyclically label a path with the ordered set of  $t$  labels  $\{x_1, \dots, x_t\}$ , label the first edge with any  $x_i$ , the second with  $x_{i+1}$ , and so on until label  $x_t$  has been used, then start with  $x_1$ , then  $x_2, \dots, x_t$ , then  $x_1$  again  $\dots$ , repeating cyclically until all edges have been labelled, with the last edge receiving label  $x_j$ . Suppose that  $N$  full cycles of labels  $\{x_1, \dots, x_t\}$  have been used. Call the matching polynomial of this labelled path  $\Phi_{i,Nt+j}$ . We show, for a fixed  $i$  and  $j$ , that  $\Phi_{i,Nt+j}$  satisfies the following recurrence, the  $(\tau, \Delta)$ -recurrence:

$$\Phi_{i,Nt+j} = \tau \Phi_{i,(N-1)t+j} - \Delta \Phi_{i,(N-2)t+j},$$

where  $\tau$  is the sum of all non-consecutive cyclic monomials in the variables  $\{x_1, \dots, x_t\}$  (see Section 1), and  $\Delta = (-1)^t x_1 \cdots x_t$ . We give two different proofs of this fact. The first one is a combinatorial/algebraic proof in Section 2 that uses the following Theorem concerning decomposing the matching polynomial  $\mathcal{M}(G, \mathbf{x})$  of a graph.

**Theorem** Let  $G$  be a labelled graph,  $H$  a subgraph of  $G$ , and  $M_H$  a matching of  $H$ , then

$$\mathcal{M}(G, \mathbf{x}) = \sum_{M_H} M_H(\mathbf{x}) \mathcal{M}(G - H - \overline{M}_H, \mathbf{x}),$$

where the summation is over every matching  $M_H$  of  $H$ . The second proof (Section 3) uses a matrix formulation of the recurrences that we develop.

Let  $G_N$  denote the first fundamental solution to the  $(\tau, \Delta)$ -recurrence; three different expressions for  $G_N$  are given in Section 4. The first expression is a sum of cyclic binomials and uses the Symmetric Representation of matrices from Section 3; the second involves Chebyshev polynomials of the second kind in the variables  $\tau$  and  $\Delta$ ; and the third is a quotient of two matching polynomials, see Theorem 4.5.

In Section 5 we extend our results from paths to cycles and rooted trees; we find explicit forms for the matching polynomial of a cyclically labelled cycle, and indicate how to find the matching polynomial of a cyclically labelled rooted tree, again using the decomposition Theorem stated above.

Many examples are given throughout the paper.

# 1 The multivariate matching polynomial of a graph, its decomposition; non-consecutive and non-consecutive cyclic functions

For a fixed  $t \geq 1$  we use multi-index notations:  $\mathbf{k} = (k_1, \dots, k_t)$ , where each  $k_s \geq 0$ ,  $\mathbf{0} = (0, \dots, 0)$ , and variables  $\mathbf{x} = (x_1, \dots, x_t)$ . The total degree of  $\mathbf{k}$  is denoted by  $|\mathbf{k}| = k_1 + \dots + k_t$ .

Let  $G$  be a finite simple graph with vertex set  $V(G)$  where  $|V(G)| \geq 1$ , and edge set  $E(G)$ . We label these edges from the  $t$  commutative variables  $\{x_1, \dots, x_t\}$ , exactly one label per edge. A *matching* of  $G$  is a collection of edges, no two of which have a vertex in common. A  $\mathbf{k}$ -*matching* of  $G$  is a matching with exactly  $k_s$  edges with label  $x_s$ , for each  $s$  with  $1 \leq s \leq t$ . If  $M_G$  is a  $\mathbf{k}$ -matching of  $G$  we define its *weight* to be

$$M_G(\mathbf{x}) = x_1^{k_1} \cdots x_t^{k_t}.$$

The empty matching of  $G$ , which contains no edges, is denoted by  $M_\emptyset$ ; it is the unique  $\mathbf{0}$ -matching and its weight is  $M_\emptyset(\mathbf{x}) = 1$ .

Define the *multivariate matching polynomial*, or simply, the *matching polynomial*, of  $G$ , by

$$\mathcal{M}(G, \mathbf{x}) = \sum_{M_G} M_G(\mathbf{x}),$$

where the summation is over every matching  $M_G$  of  $G$ .

Denote the number of  $\mathbf{k}$ -matchings of  $G$  by  $a(G, \mathbf{k})$ . Then an alternative definition of the multivariate matching polynomial of  $G$  is

$$\mathcal{M}(G, \mathbf{x}) = \sum_{(k_1, \dots, k_t)} a(G, \mathbf{k}) x_1^{k_1} \cdots x_t^{k_t}.$$

The multivariate matching polynomial is a natural extension of the matching polynomial of Farrell [1]. Indeed, here with  $t = 1$  and in [1] with  $w_1 = 1$  and  $w_2 = x_1$ , the polynomials are identical.

Let  $P_1$  be the graph with one vertex and no edges, *i.e.*, an isolated vertex; we define  $\mathcal{M}(P_1, \mathbf{x}) = 1$ . Now suppose  $G' = G \cup nP_1$ , where  $n \geq 1$ , *i.e.*,  $G'$

is the disjoint union of  $G$  and  $n$  isolated vertices, then we define  $\mathcal{M}(G', \mathbf{x}) = \mathcal{M}(G, \mathbf{x})$ .

For any edge  $e \in E(G)$ , let  $\bar{e}$  denote the set of edges that are incident to  $e$ ; and for any subgraph  $H$  of  $G$ , let  $\bar{H} = \cup_{e \in E(H)} \bar{e}$ . Define  $\bar{M}_\emptyset = \emptyset$ . Also let  $G - H$  be the graph obtained from  $G$  when all the edges of  $H$  are removed, so  $G - H$  has the same vertex set as  $G$ .

Now let  $H$  be a fixed subgraph of  $G$  and let  $M_H$  be a matching of  $H$ . In the following theorem we express  $\mathcal{M}(G, \mathbf{x})$  as a sum of terms, each term containing the weight of a fixed matching,  $M_H(\mathbf{x})$ , of  $H$ ; we call this *decomposing*  $\mathcal{M}(G, \mathbf{x})$  at  $H$ .

**Theorem 1.1** *Let  $G$  be a graph labelled as above,  $H$  a fixed subgraph of  $G$ , and  $M_H$  a matching of  $H$ . Then*

$$\mathcal{M}(G, \mathbf{x}) = \sum_{M_H} M_H(\mathbf{x}) \mathcal{M}(G - H - \bar{M}_H, \mathbf{x}), \quad (1)$$

where the summation is over every matching  $M_H$  of  $H$ .

*Proof.* Let  $M_G$  be a matching of  $G$  which induces a (fixed) matching  $M_H$  on  $H$ , i.e.,  $M_G$  contains exactly  $M_H$  and no other edges from  $H$ . Then  $M_G(\mathbf{x}) = M_H(\mathbf{x}) M(\mathbf{x})$  where  $M$  is a matching of  $G$  with no edges in  $H$ , and also with no edges in  $\bar{M}_H$  or else  $M_G$  would not be a matching. Hence,  $M$  is a matching of  $G - H - \bar{M}_H$ , i.e.,  $M(\mathbf{x})$  is a term of  $\mathcal{M}(G - H - \bar{M}_H, \mathbf{x})$ . So  $M_H(\mathbf{x}) \mathcal{M}(G - H - \bar{M}_H, \mathbf{x})$  is the sum of the weights of all the matchings in  $G$  which induce  $M_H$  on  $H$ .

Now every matching in  $G$  induces some matching on  $H$ , so we may sum over all matchings in  $H$  to give (1). ■

Theorem 1.1 extends known facts about matching polynomials, eg., see Theorem 1 of Farrell [1] for the case where  $H$  is a single edge. We have the corresponding:

**Corollary 1.2** *Let  $G$  be a graph labelled as above, and let  $H = e$  labelled with  $x$  be an edge of  $G$ . Then*

$$\mathcal{M}(G, \mathbf{x}) = \mathcal{M}(G - e, \mathbf{x}) + x \mathcal{M}(G - e - \bar{e}, \mathbf{x}). \quad (2)$$

*Proof.* The result comes from (1) since  $H = e$  has just two matchings: the empty matching  $M_\emptyset$  with weight  $M_\emptyset(\mathbf{x}) = 1$ , and the matching  $e$  with weight  $M_e(\mathbf{x}) = x$ .  $\blacksquare$

**Notation** Throughout this paper we use  $P_m$  to denote the path with  $m$  vertices and  $m - 1$  edges.

Fix  $i$  and  $j$  where  $1 \leq i \leq j \leq t$ . Consider the path  $P_{j-i+2}$  with its  $j-i+1$  edges labelled from the ordered set  $\{x_i, \dots, x_j\}$ , the first edge receiving label  $x_i$ , and the last  $x_j$ ; see Fig. 1.




Fig. 1: The labelled path  $P_{j-i+2}$  with matching polynomial  $\phi_{i,j}$ .

The pair  $x_s x_{s+1}$  for any fixed  $s$  with  $i \leq s \leq j - 1$  is called a *consecutive* pair. A monomial from the ordered set  $\{x_i, \dots, x_j\}$  that contains no consecutive pairs is a *non-consecutive monomial*, a *nc-monomial*. Note that the empty monomial is a *nc-monomial* that we denote by 1.

Let  $\phi_{i,j}$  be the sum of all *nc-monomials* in the ordered variables  $\{x_i, \dots, x_j\}$ . Then  $\phi_{i,j} = \mathcal{M}(P_{j-i+2}, \mathbf{x})$  is the matching polynomial of the labelled path  $P_{j-i+2}$ . We call the functions  $\phi_{i,j}$  *elementary non-consecutive functions*, and for any  $i \geq 1$  define the initial values

$$\phi_{i,i-2} = \phi_{i,i-1} = 1. \quad (3)$$

These initial values ensure that the following recurrence is valid for any  $j$  with  $i \leq j \leq t$ .

**Theorem 1.3** *For a fixed  $i$  and  $j$  with  $1 \leq i \leq j \leq t$  and the initial values in (3), we have*

$$\phi_{i,j} = \phi_{i,j-1} + x_j \phi_{i,j-2}. \quad (4)$$

*Proof.* Let  $e$  be the rightmost edge of  $G = P_{j-i+2}$  shown in Fig. 1, and apply (2).  $\blacksquare$

**Example 1** For arbitrary  $i$  we have

$$\begin{aligned}\phi_{i,i} &= 1 + x_i, & \phi_{i,i+1} &= 1 + x_i + x_{i+1}, \\ \phi_{i,i+2} &= 1 + x_i + x_{i+1} + x_{i+2} + x_i x_{i+2}, \\ \phi_{i,i+3} &= 1 + x_i + x_{i+1} + x_{i+2} + x_{i+3} + x_i x_{i+2} + x_i x_{i+3} + x_{i+1} x_{i+3}.\end{aligned}$$

**Example 2** For arbitrary  $i$ , putting  $j = i - 1$  and  $j = i - 2$  in Recurrence (4) and using (3) give

$$\phi_{i,i-3} = 0 \quad \text{and} \quad \phi_{i,i-4} = \frac{1}{x_{i-2}}.$$

In the second equation, if  $i = 1$  we replace  $x_{-1}$  by  $x_{t-1}$ , and if  $i = 2$  we replace  $x_0$  by  $x_t$ .

Consider Recurrence (4). It is convenient to work with a basis of solutions to this recurrence. Denote the first fundamental solution by  $f_{i,j}$  and the second by  $g_{i,j}$ , with initial values

$$f_{i,i-2} = 0, f_{i,i-1} = 1 \quad \text{and} \quad g_{i,i-2} = 1, g_{i,i-1} = 0. \quad (5)$$

So

$$\phi_{i,i-2} = f_{i,i-2} + g_{i,i-2} \quad \text{and} \quad \phi_{i,i-1} = f_{i,i-1} + g_{i,i-1}.$$

Now, from Recurrence (4) and strong induction on  $j$ , we have (6) below for all  $j$  with  $i \leq j \leq t$

$$\phi_{i,j} = f_{i,j} + g_{i,j}. \quad (6)$$

$$\phi_{i,j} = \phi_{i+1,j} + x_i \phi_{i+2,j}. \quad (7)$$

Equation (7) comes from decomposing  $\phi_{i,j}$  at the leftmost edge of  $P_{j-i+2}$ , whose label is  $x_i$ , *i.e.*, decomposing  $\phi_{i,j}$  at  $x_i$ ; see Corollary 1.2. These two equations suggest that the fundamental solutions are given by

$$f_{i,j} = \phi_{i+1,j} \quad \text{and} \quad g_{i,j} = x_i \phi_{i+2,j}.$$

This is indeed the case:

**Lemma 1.4** *For any  $j$  with  $i \leq j \leq t$  we have*

- (i)  $f_{i,j} = \phi_{i+1,j}$ ,
- (ii)  $g_{i,j} = x_i \phi_{i+2,j}$ .

*Proof.* We need only prove (i) because of (6) and (7) above.

From (5) we have  $f_{i,i-2} = 0$  and from Example 2 we have  $\phi_{i+1,i-2} = 0$ ; thus  $f_{i,i-2} = \phi_{i+1,i-2}$ . Similarly, from (5) and (3), we have  $f_{i,i-1} = \phi_{i+1,i-1}$ . So both  $f_{i,j}$  and  $\phi_{i+1,j}$  have the same initial values at  $j = i - 2$  and  $j = i - 1$  and they both satisfy Recurrence (4), so they are equal for any  $j$  with  $i \leq j \leq t$ . ■

Thus we know combinatorially what the two fundamental solutions to Recurrence (4) are. The first,  $f_{i,j}$ , is the matching polynomial of the path shown in Fig. 2(a); the second,  $g_{i,j}$ , is  $x_i \times$  the matching polynomial of the path in Fig. 2(b).




Fig. 2 (a) The labelled path with matching polynomial  $f_{i,j}$ .

(b) The labelled path with matching polynomial  $\frac{g_{i,j}}{x_i}$ .

**Example 3** For arbitrary  $i$  we have

$$\begin{aligned} f_{i,i} &= 1, & g_{i,i} &= x_i, \\ f_{i,i+1} &= 1 + x_{i+1}, & g_{i,i+1} &= x_i, \\ f_{i,i+2} &= 1 + x_{i+1} + x_{i+2}, & g_{i,i+2} &= x_i + x_i x_{i+2}. \end{aligned}$$

Now arrange the variables  $\{x_i, \dots, x_j\}$  clockwise around a circle. Thus  $x_i$  and  $x_j$  are consecutive. Call a pair  $x_s x_{s'}$  *consecutive cyclic* if  $x_s$  and  $x_{s'}$  are consecutive on this circle. Call a monomial from  $\{x_i, \dots, x_j\}$  a *non-consecutive cyclic* monomial — *ncc*-monomial — if it contains no consecutive cyclic pairs. The empty monomial is a *ncc*-monomial that we denote by 1.

Let  $\tau_{i,j}$  be the sum of all *ncc*-monomials in the variables  $\{x_i, \dots, x_j\}$ . Then, for  $j \geq i + 2$ ,  $\tau_{i,j} = \mathcal{M}(C_{j-i+1}, \mathbf{x})$  is the matching polynomial of the



labelled cycle  $C_{j-i+1}$  with  $j - i + 1$  edges and  $j - i + 1$  vertices, shown in Fig. 3; the cycle starts at the large vertex, and proceeds clockwise.




Fig. 3: The labelled cycle  $C_{j-i+1}$  with matching polynomial  $\tau_{i,j}$ .

For initial values let

$$\tau_{i,i-1} = 2, \quad \tau_{i,i} = 1, \quad \text{and} \quad \tau_{i,i+1} = 1 + x_i + x_{i+1}. \quad (8)$$

**Lemma 1.5** *For any  $j$  with  $i \leq j \leq t$  we have*

- (i)  $\tau_{i,j} = f_{i,j} + g_{i,j-1}$ ,
- (ii)  $\phi_{i,j} - \tau_{i,j} = x_i x_j \phi_{i+2,j-2}$ .

*Proof.* (i) We check this equality at  $j = i$  and  $j = i + 1$  using (5), Example 3, and (8). For  $j \geq i + 2$  we decompose  $\tau_{i,j}$  at  $x_i$  yielding  $\tau_{i,j} = \phi_{i+1,j} + x_i \phi_{i+2,j-1}$ , which gives (i) via Lemma 1.4.

(ii) We check at  $j = i$  and  $j = i + 1$  using Examples 1 and 2, and (8). For  $j \geq i + 2$  the difference  $\phi_{i,j} - \tau_{i,j}$  consists of all  $nc$ -monomials that contain the consecutive cyclic pair  $x_i x_j$ ; clearly this is  $x_i x_j \times$  the sum of all  $nc$ -monomials on  $\{x_{i+2}, \dots, x_{j-2}\}$ , *i.e.*,  $x_i x_j \phi_{i+2,j-2}$ . ■

**Example 4** For arbitrary  $i$  we have

$$\begin{aligned} \tau_{i,i+2} &= 1 + x_i + x_{i+1} + x_{i+2}, \\ \tau_{i,i+3} &= 1 + x_i + x_{i+1} + x_{i+2} + x_{i+3} + x_i x_{i+2} + x_{i+1} x_{i+3}. \end{aligned}$$

## 2 Cyclically labelled paths; $\Phi_{i,Nt+j}$ and the $(\tau, \Delta)$ -recurrence

Consider a path  $P$  and the ordered set of  $t$  labels  $\{x_1, \dots, x_t\}$ . For a fixed  $i$ , where  $1 \leq i \leq t$ , and moving from left to right, label the first edge of  $P$  with  $x_i$ , the second with  $x_{i+1}$ , and so on until label  $x_t$  has been used; so the  $(t-i+1)$ -th edge receives label  $x_t$ . Then label edge  $t-i+2$  with  $x_1$ , and edge  $t-i+3$  with  $x_2$ , and so on  $\dots$ , labelling cyclically with  $\{x_1, \dots, x_t\}$  until all edges have been labelled. Let the last edge receive label  $x_j$ , where  $1 \leq j \leq t$ . Suppose that  $N \geq 0$  full cycles of labels  $\{x_1, \dots, x_t\}$  have been used beginning at edge  $t-i+2$ . Then if  $j = t$  we call this path  $P(i, Nt)$ , or if  $1 \leq j < t$  we call it  $P(i, Nt+j)$ . This labelling is a *cyclic labelling*. The cyclically labelled path  $P(i, Nt+j)$  is shown in Fig. 4. Let  $\Phi_{i,Nt+j}(\mathbf{x}) = \Phi_{i,Nt+j}$  denote the matching polynomial of the path  $P(i, Nt+j)$ .




Fig. 4: The cyclically labelled path  $P(i, Nt+j)$  with matching polynomial  $\Phi_{i,Nt+j}$ .

We define the initial conditions for  $\Phi_{i,Nt+j}$  as

$$\begin{aligned} N = -1 : \quad & \Phi_{i,-t+j} = \phi_{i,j}, \quad \text{for all } j \text{ with } 0 \leq j \leq t, \\ \text{also } N = 0 : \quad & \Phi_{i,0t+j} = \Phi_{i,j}. \end{aligned} \quad (9)$$

In order to find  $\phi_{i,j}$  if  $j < i$  we use the initial values for  $\phi_{i,j}$  from (3) and push back Recurrence (4), as shown in Example 2.

Now  $\Phi_{i,Nt+j}$  satisfies the same recurrence as that of  $\phi_{i,j}$ , Recurrence (4); the proof is similar, noting that  $x_0$  must be replaced by  $x_t$ , and considering  $Nt-1$  as  $(N-1)t+t-1$ , etc.

**Lemma 2.1** *For any  $N \geq -1$  and  $j$  with  $0 \leq j \leq t$  we have*

$$\Phi_{i,Nt+j} = \Phi_{i,Nt+j-1} + x_j \Phi_{i,Nt+j-2}. \quad (10)$$

■

**Notation** For  $i = 1$  we write  $\phi_{i,j} = \phi_{1,j} = \phi_j$  and  $\phi_t = \phi$ , also  $\tau_{1,j} = \tau_j$  and  $\tau_t = \tau$ , and  $f_{1,j} = f_j$ , etc. Also let  $\Delta = (-1)^t x_1 \cdots x_t$ .

**Lemma 2.2** For any  $N \geq 0$  and any  $j$  with  $0 \leq j \leq t$  we have

$$\Phi_{i,Nt+j} = \Phi_{i,Nt} f_j + \Phi_{i,Nt-1} g_j. \quad (11)$$

*Proof.* With  $N = 0$  and  $j = 0$  Equation (11) is true using the initial values  $f_0 = 1$  and  $g_0 = 0$  of (5) with  $i = 1$ . Otherwise, consider the path  $P(i, Nt + j)$  of Fig. 4 and decompose its matching polynomial,  $\Phi_{i,Nt+j}$ , at the edge labelled  $x_1$  marked with a  $*$ . This gives

$$\begin{aligned} \Phi_{i,Nt+j} &= \Phi_{i,Nt} \phi_{2,j} + x_1 \Phi_{i,Nt-1} \phi_{3,j} \\ &= \Phi_{i,Nt} f_j + \Phi_{i,Nt-1} g_j, \end{aligned}$$

using Lemma 1.4. ■

Now we define the second order  $(\tau, \Delta)$ -recurrence

$$\Theta_N = \tau \Theta_{N-1} - \Delta \Theta_{N-2}. \quad (12)$$

Let  $G_N(\mathbf{x}) = G_N$  denote the first fundamental solution to this recurrence. We will evaluate  $G_N$  in Section 4.

In Theorem 2.4 below we show that, for a fixed  $i$  and  $j$ ,  $\Phi_{i,Nt+j}$  satisfies the  $(\tau, \Delta)$ -recurrence. First:

**Lemma 2.3** For any  $N \geq 1$  we have

$$\begin{aligned} (i) \quad &\Phi_{i,Nt-1} f_t - \Phi_{i,Nt} f_{t-1} = \Delta \Phi_{i,(N-1)t-1}, \\ (ii) \quad &\Phi_{i,Nt-1} g_t - \Phi_{i,Nt} g_{t-1} = -\Delta \Phi_{i,(N-1)t}. \end{aligned} \quad (13)$$

*Proof.* (i) Using Recurrence (4) on  $f_t$  and on  $\Phi_{i,Nt}$  (see Lemma 2.1), the left-hand side of (13) becomes

$$\Phi_{i,Nt-1} \{f_{t-1} + x_t f_{t-2}\} - \{\Phi_{i,Nt-1} + x_t \Phi_{i,Nt-2}\} f_{t-1} = -x_t \{\Phi_{i,Nt-2} f_{t-1} - \Phi_{i,Nt-1} f_{t-2}\}.$$

The second factor in the right-hand side of this equation is the left-hand side of (13) with subscripts shifted down by 1. After  $t$  such iterations the left-hand side of (13) becomes

$$(-x_t)(-x_{t-1}) \cdots (-x_1) \{\Phi_{i,(N-1)t-1} f_0 - \Phi_{i,(N-1)t} f_{-1}\} = \Delta \Phi_{i,(N-1)t-1},$$

using the initial values  $f_0 = 1$  and  $f_{-1} = 0$ . The proof of (ii) is similar. ■

Now a main result:  $\Phi_{i,Nt+j}$  satisfies the  $(\tau, \Delta)$ -recurrence.

**Theorem 2.4** *For any  $N \geq 1$ , and any fixed  $i$  with  $1 \leq i \leq t$ , and any fixed  $j$  with  $0 \leq j \leq t$ , we have*

$$\Phi_{i,Nt+j} = \tau \Phi_{i,(N-1)t+j} - \Delta \Phi_{i,(N-2)t+j}. \quad (14)$$

*Proof.* Due to Recurrences (4) and (10) we need only show that (14) is true when  $j = t$  and  $t - 1$ . It will then be true for all  $j$  with  $0 \leq j \leq t$  by pushing back Recurrence (10).

With  $N \geq 1$  and  $j = t$ , Equation (11) gives

$$\begin{aligned} \Phi_{i,Nt+t} &= \Phi_{i,Nt} f_t + \Phi_{i,Nt-1} g_t \\ &= \Phi_{i,Nt} f_t + \Phi_{i,Nt-1} g_t + \Phi_{i,Nt} g_{t-1} - \Phi_{i,Nt} g_{t-1} \\ &= \Phi_{i,Nt} f_t + \Phi_{i,Nt} g_{t-1} + \Phi_{i,Nt-1} g_t - \Phi_{i,Nt} g_{t-1} \\ &= \tau \Phi_{i,Nt} - \Delta \Phi_{i,(N-1)t}, \\ &= \tau \Phi_{i,(N-1)t+t} - \Delta \Phi_{i,(N-2)t+t}, \end{aligned}$$

using  $\tau = \tau_t = f_t + g_{t-1}$  from Lemma 1.5(i), and Lemma 2.3(ii) at the fourth line. For  $j = t - 1$  the proof is similar using Lemma 2.3(i). ■

### 3 Matrix formulation of recurrences

Here we use matrices to give another proof that  $\Phi_{i,Nt+j}$  satisfies the  $(\tau, \Delta)$ -recurrence, and prepare for the evaluation of  $G_N$  in Section 4.

Recall from Section 1 that  $f_{i,j}$  and  $g_{i,j}$  are the 2 fundamental solutions to Recurrence (4). Now define the matrix

$$X_{i,j} = \begin{pmatrix} g_{i,j-1} & f_{i,j-1} \\ g_{i,j} & f_{i,j} \end{pmatrix}.$$

Then the recurrences for  $f_{i,j}$  and  $g_{i,j}$  can be written as:

$$X_{i,j} = \begin{pmatrix} 0 & 1 \\ x_j & 1 \end{pmatrix} \begin{pmatrix} g_{i,j-2} & f_{i,j-2} \\ g_{i,j-1} & f_{i,j-1} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ x_j & 1 \end{pmatrix} X_{i,j-1}. \quad (15)$$

Consistent with (5) we have  $X_{i,i-1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = I$ , the  $2 \times 2$  identity matrix.

Thus, for  $j \geq i$ , we have

$$X_{i,j} = \begin{pmatrix} 0 & 1 \\ x_j & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ x_{j-1} & 1 \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 \\ x_i & 1 \end{pmatrix}. \quad (16)$$

Let  $\mathbf{1} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$  and  $\mathbf{e} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ , and let  $\langle \cdot, \cdot \rangle$  denote the usual inner product. Then for  $j \geq i$ , and using (6),

$$\phi_{i,j} = \langle X_{i,j} \mathbf{1}, \mathbf{e} \rangle. \quad (17)$$

As before if  $i = 1$  we let  $X_{1,j} = X_j$  and if  $j = t$  we let  $X = X_t$ , in particular,

$$X = \begin{pmatrix} g_{t-1} & f_{t-1} \\ g_t & f_t \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ x_t & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ x_{t-1} & 1 \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 \\ x_1 & 1 \end{pmatrix}. \quad (18)$$

For  $N \geq 0$ , from (10) we may also write

$$\begin{pmatrix} \Phi_{i,Nt+j-1} \\ \Phi_{i,Nt+j} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ x_j & 1 \end{pmatrix} \begin{pmatrix} \Phi_{i,Nt+j-2} \\ \Phi_{i,Nt+j-1} \end{pmatrix},$$

and then repeated use of (15) gives

$$\Phi_{i,Nt+j} = \langle X_j X^N X_{i,t} \mathbf{1}, \mathbf{e} \rangle. \quad (19)$$

Now using (16) and (18) we see that  $X_j X^{-1} X_{i,t} = X_{i,j}$ . So, using (17) and (9), we have

$$\langle X_j X^{-1} X_{i,t} \mathbf{1}, \mathbf{e} \rangle = \langle X_{i,j} \mathbf{1}, \mathbf{e} \rangle = \phi_{i,j} = \Phi_{i,-t+j},$$

thus (19) is true for  $N = -1$  also.

**Theorem 3.1** For  $N \geq -1$  we have

$$\Phi_{i,Nt+j} = \langle X_j X^N X_{i,t} \mathbf{1}, \mathbf{e} \rangle.$$

■

From Lemma 1.5(i) and (16) we have the following forms for the trace and determinant of matrix  $X_{i,j}$

$$\text{tr}(X_{i,j}) = \tau_{i,j} \quad \text{and} \quad \det(X_{i,j}) = (-1)^{j-i+1} x_i \cdots x_j.$$

In particular, for matrix  $X$  from (18), we have

$$\text{tr}(X) = \tau \quad \text{and} \quad \det(X) = \Delta. \quad (20)$$

Now let  $Z$  be any invertible  $2 \times 2$  matrix with trace  $\text{tr}(Z)$  and determinant  $\det(Z)$ , and let  $T$  denote transpose. Then the Cayley-Hamilton theorem says that  $Z^2 = \text{tr}(Z) Z - \det(Z) I$ , so  $Z^N = \text{tr}(Z) Z^{N-1} - \det(Z) Z^{N-2}$ , for  $N \geq 1$ . Let  $\mathbf{u}$  and  $\mathbf{v} \in \mathbf{R}^2$  and, for  $N \geq -1$ , define  $\Psi_N = \langle Z^N \mathbf{u}, \mathbf{v} \rangle$ . Then

**Lemma 3.2** For  $N \geq 1$ ,  $\Psi_N$  satisfies the recurrence

$$\Psi_N = \text{tr}(Z) \Psi_{N-1} - \det(Z) \Psi_{N-2},$$

with initial conditions  $\Psi_{-1} = \langle Z^{-1} \mathbf{u}, \mathbf{v} \rangle$  and  $\Psi_0 = \langle Z^0 \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{u}, \mathbf{v} \rangle$ .

■

Now for  $N \geq -1$ ,

$$\Phi_{i,Nt+j} = \langle X_j X^N X_{i,t} \mathbf{1}, \mathbf{e} \rangle = \langle X^N X_{i,t} \mathbf{1}, X_j^T \mathbf{e} \rangle.$$

So, for  $N \geq 1$ , Lemma 3.2 with  $Z = X$ ,  $\mathbf{u} = X_{i,t} \mathbf{1}$ , and  $\mathbf{v} = X_j^T \mathbf{e}$ , and (20), gives,

$$\Phi_{i,Nt+j} = \tau \Phi_{i,(N-1)t+j} - \Delta \Phi_{i,(N-2)t+j}.$$

This is a second proof that  $\Phi_{i,Nt+j}$  satisfies the  $(\tau, \Delta)$ -recurrence.

## 4 The Symmetric Representation, MacMahon's Master Theorem, three expressions for $G_N$

Consider polynomials in the variables  $u_1, \dots, u_d$ . We will work with the vector space whose basis elements are the homogeneous polynomials of degree  $N$  in these variables, *i.e.*, with

$$\{u_1^{n_1} \cdots u_d^{n_d} \mid n_1 + \cdots + n_d = N, \text{ each } n_\ell \geq 0\},$$

this vector space has dimension  $\binom{N+d-1}{N}$ .

The symmetric representation of a  $d \times d$  matrix  $A = (a_{\ell\ell'})$  is the action on polynomials induced by:

$$u_1^{n_1} \cdots u_d^{n_d} \rightarrow v_1^{n_1} \cdots v_d^{n_d},$$

where

$$v_\ell = \sum_{\ell'} a_{\ell\ell'} u_{\ell'}$$

or, more compactly,  $v = Au$ . That is, define the matrix element  $\left\langle \begin{matrix} m_1, \dots, m_d \\ n_1, \dots, n_d \end{matrix} \right\rangle_A$  to be the coefficient of  $u_1^{n_1} \cdots u_d^{n_d}$  in  $v_1^{m_1} \cdots v_d^{m_d}$ . Then, for a fixed  $(m_1, \dots, m_d)$ , we have

$$v_1^{m_1} \cdots v_d^{m_d} = \sum_{(n_1, \dots, n_d)} \left\langle \begin{matrix} m_1, \dots, m_d \\ n_1, \dots, n_d \end{matrix} \right\rangle_A u_1^{n_1} \cdots u_d^{n_d}. \quad (21)$$

Observe that the total degree  $N = |n| = \sum n_\ell = |m| = \sum m_\ell$ , *i.e.*, homogeneity of degree  $N$  is preserved. We use multi-indices:  $m = (m_1, \dots, m_d)$  and  $n = (n_1, \dots, n_d)$ . Then, for a fixed  $m$ , (21) becomes

$$v^m = \sum_n \left\langle \begin{matrix} m \\ n \end{matrix} \right\rangle_A u^n.$$

Successive application of  $B$  then  $A$  shows that this is a homomorphism of the multiplicative semi-group of square  $d \times d$  matrices into the multiplicative semi-group of square  $\binom{N+d-1}{N} \times \binom{N+d-1}{N}$  matrices.

**Proposition 4.1** *Matrix elements satisfy the homomorphism property*

$$\left\langle \begin{matrix} m \\ n \end{matrix} \right\rangle_{AB} = \sum_k \left\langle \begin{matrix} m \\ k \end{matrix} \right\rangle_A \left\langle \begin{matrix} k \\ n \end{matrix} \right\rangle_B.$$

*Proof.* Let  $v = (AB)u$  and  $w = Bu$ . Then,

$$\begin{aligned} v^m &= \sum_n \left\langle \begin{matrix} m \\ n \end{matrix} \right\rangle_{AB} u^n \\ &= \sum_k \left\langle \begin{matrix} m \\ k \end{matrix} \right\rangle_A w^k \\ &= \sum_n \sum_k \left\langle \begin{matrix} m \\ k \end{matrix} \right\rangle_A \left\langle \begin{matrix} k \\ n \end{matrix} \right\rangle_B u^n. \end{aligned}$$

■

**Definition** Fix the degree  $N = \sum n_\ell = \sum m_\ell$ . Define  $\text{tr}_{\text{Sym}}^N(A)$ , the *symmetric trace* of  $A$  in degree  $N$ , as the sum of the diagonal elements  $\left\langle \begin{matrix} m \\ n \end{matrix} \right\rangle_A$ , i.e.,

$$\text{tr}_{\text{Sym}}^N(A) = \sum_m \left\langle \begin{matrix} m \\ m \end{matrix} \right\rangle_A.$$

Equality such as  $\text{tr}_{\text{Sym}}(A) = \text{tr}_{\text{Sym}}(B)$  means that the symmetric traces are equal in every degree  $N \geq 0$ .

**Remark** The action defined here on polynomials is equivalent to the action on symmetric tensor powers, see Fulton and Harris [2], pp. 472-5.

Now it is straightforward to see directly (cf. the diagonal case shown in the Corollary below) that if  $A$  is upper-triangular, with eigenvalues  $\lambda_1, \dots, \lambda_d$ , then  $\text{tr}_{\text{Sym}}^N(A) = h_N(\lambda_1, \dots, \lambda_d)$ , the  $N^{\text{th}}$  homogeneous symmetric function. The homomorphism property, Proposition 4.1, shows that  $\text{tr}_{\text{Sym}}^N(AB) = \text{tr}_{\text{Sym}}^N(BA)$ , and that similar matrices have the same trace. Again by the homomorphism property, if two  $d \times d$  matrices are similar,  $A = MBM^{-1}$ , then that relation extends to their respective symmetric representations in every degree. Recall that any matrix is similar to an upper-triangular one with the same eigenvalues. Thus,

**Theorem 4.2** Symmetric Trace Theorem (see pp. 51-2 of Springer [5]).

*We have*

$$\frac{1}{\det(I - cA)} = \sum_{N=0}^{\infty} c^N \text{tr}_{\text{Sym}}^N(A).$$

*Proof.* With  $\lambda_\ell$  denoting the eigenvalues of  $A$ ,

$$\frac{1}{\det(I - cA)} = \prod_{\ell} \frac{1}{1 - c\lambda_\ell}$$



$$\begin{aligned}
&= \sum_{N=0}^{\infty} c^N h_N(\lambda_1, \dots, \lambda_d) \\
&= \sum_{N=0}^{\infty} c^N \text{tr}_{\text{Sym}}^N(A).
\end{aligned}$$

■

As a Corollary we have MacMahon's Master Theorem, which we express in the above terminology.

**Corollary 4.3** MacMahon's Master Theorem.

The diagonal matrix element  $\left\langle \begin{smallmatrix} m \\ m \end{smallmatrix} \right\rangle_A$  is the coefficient of  $u^m = u_1^{m_1} \dots u_d^{m_d}$  in the expansion of  $\det(I - UA)^{-1}$  where  $U = \text{diag}(u_1, \dots, u_d)$  is the diagonal matrix with entries  $u_1, \dots, u_d$  on the diagonal.

*Proof.* From Theorem 4.2, with  $c = 1$ , we want to calculate the symmetric trace of  $UA$ . By the homomorphism property,

$$\begin{aligned}
\text{tr}_{\text{Sym}}^N(UA) &= \sum_m \left\langle \begin{smallmatrix} m \\ m \end{smallmatrix} \right\rangle_{UA} \\
&= \sum_m \sum_k \left\langle \begin{smallmatrix} m \\ k \end{smallmatrix} \right\rangle_U \left\langle \begin{smallmatrix} k \\ m \end{smallmatrix} \right\rangle_A.
\end{aligned}$$

Now, with  $v = Uw$  and  $v_\ell = u_\ell w_\ell$ , then

$$v^m = (u_1 w_1)^{m_1} \dots (u_d w_d)^{m_d} = u^m w^m = \sum_k \left\langle \begin{smallmatrix} m \\ k \end{smallmatrix} \right\rangle_U w^k,$$

*i.e.*,

$$\left\langle \begin{smallmatrix} m \\ k \end{smallmatrix} \right\rangle_U = u_1^{m_1} \dots u_d^{m_d} \delta_{k_1 m_1} \dots \delta_{k_d m_d}$$

so that

$$\text{tr}_{\text{Sym}}^N(UA) = \sum_m \left\langle \begin{smallmatrix} m \\ m \end{smallmatrix} \right\rangle_A u^m.$$

■

Now we restrict ourselves to  $d = 2$ , and return to the  $(\tau, \Delta)$ -recurrence.

Recall, from (18), the  $2 \times 2$  matrix

$$\begin{aligned} X &= \begin{pmatrix} 0 & 1 \\ x_t & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ x_{t-1} & 1 \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 \\ x_1 & 1 \end{pmatrix} \\ &= \xi_t \xi_{t-1} \cdots \xi_1, \end{aligned}$$

where  $\xi_s = \begin{pmatrix} 0 & 1 \\ x_s & 1 \end{pmatrix}$  for  $1 \leq s \leq t$ . Let us modify  $\xi_s$  slightly by defining  $\alpha_s = \begin{pmatrix} 0 & 1 \\ x_s & a_s \end{pmatrix}$  for  $1 \leq s \leq t$ , and calling

$$\begin{aligned} \bar{X} &= \begin{pmatrix} 0 & 1 \\ x_t & a_t \end{pmatrix} \begin{pmatrix} 0 & 1 \\ x_{t-1} & a_{t-1} \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 \\ x_1 & a_1 \end{pmatrix} \\ &= \alpha_t \alpha_{t-1} \cdots \alpha_1. \end{aligned}$$

Let

$$\text{tr}(\bar{X}) = \bar{\tau} \quad \text{and} \quad \det(\bar{X}) = \bar{\Delta},$$

and let  $\bar{G}_N$  be the first fundamental solution to the  $(\bar{\tau}, \bar{\Delta})$ -recurrence:

$$\Theta_N = \bar{\tau} \Theta_{N-1} - \bar{\Delta} \Theta_{N-2}. \quad (22)$$

Then

$$\begin{aligned} \sum_{N=0}^{\infty} c^N \bar{G}_N &= \frac{1}{1 - \bar{\tau}c + \bar{\Delta}c^2} \\ &= \frac{1}{\det(I - c\bar{X})} \\ &= \sum_{N=0}^{\infty} c^N \text{tr}_{\text{Sym}}^N(\bar{X}). \end{aligned}$$

So

$$\bar{G}_N = \text{tr}_{\text{Sym}}^N(\bar{X}) = \sum_m \langle m \rangle_{\bar{X}} = \sum_m \langle m \rangle_{\alpha_t \alpha_{t-1} \cdots \alpha_1}.$$

We need to calculate the symmetric trace of  $\bar{X}$  and so identify  $\bar{G}_N$ . By the homomorphism property, we need only find the matrix elements for each matrix  $\alpha_s$ , multiply together and take the trace.

For  $\alpha_s = \begin{pmatrix} 0 & 1 \\ x_s & a_s \end{pmatrix}$  the mapping induced on polynomials is

$$v_1 = u_2, \quad v_2 = x_s u_1 + a_s u_2. \quad (23)$$

For any integer  $N \geq 0$ , the expansion of  $v_1^m v_2^{N-m}$  in powers of  $u_1$  and  $u_2$  is of the form

$$v_1^m v_2^{N-m} = \sum_n \langle m \rangle_n \alpha_s u_1^n u_2^{N-n}, \quad (24)$$

with the notation for the matrix elements abbreviated accordingly. From (23) and (24), the binomial theorem yields

$$\langle m \rangle_n \alpha_s = \binom{N-m}{n} x_s^n a_s^{N-m-n}.$$

For example, when  $t = 3$ , the product  $\bar{X} = \alpha_3 \alpha_2 \alpha_1$  gives the matrix elements, for homogeneity of degree  $N$ ,

$$\begin{aligned} \langle m \rangle_n \bar{X} &= \sum_{(k_2, k_3)} \langle m \rangle_{k_3} \alpha_3 \langle k_3 \rangle_{k_2} \alpha_2 \langle k_2 \rangle_n \alpha_1 \\ &= \sum_{(k_2, k_3)} \binom{N-m}{k_3} \binom{N-k_3}{k_2} \binom{N-k_2}{n} x_1^n x_2^{k_2} x_3^{k_3} a_1^{N-k_2-n} a_2^{N-k_3-k_2} a_3^{N-k_3-m}. \end{aligned}$$

Thus, the symmetric trace  $\text{tr}_{\text{Sym}}^N(\bar{X}) = \sum_m \langle m \rangle_m \bar{X}$  is

$$\sum_{(k_1, k_2, k_3)} \binom{N-k_2}{k_1} \binom{N-k_3}{k_2} \binom{N-k_1}{k_3} x_1^{k_1} x_2^{k_2} x_3^{k_3} a_1^{N-k_1-k_2} a_2^{N-k_2-k_3} a_3^{N-k_3-k_1},$$

a *cyclic binomial*. In general, for a product of arbitrary length, the symmetric trace is given by the corresponding cyclic binomial.

Recall the recurrence

$$S_N(x) = 2x S_{N-1}(x) - S_{N-2}(x), \quad (25)$$

for  $N \geq 1$ . The Chebyshev polynomials of the first kind,  $T_N = T_N(x)$ , are solutions of this recurrence with initial conditions  $T_{-1} = x$  and  $T_0 = 1$ , and the Chebyshev polynomials of the second kind,  $U_N = U_N(x)$ , are solutions with  $U_{-1} = 0$  and  $U_0 = 1$ .

Combining these observations yields the main identities:

**Theorem 4.4** Let  $\overline{X} = \alpha_t \alpha_{t-1} \cdots \alpha_1$ , with  $\alpha_s = \begin{pmatrix} 0 & 1 \\ x_s & a_s \end{pmatrix}$  for  $1 \leq s \leq t$ , and let  $\overline{\tau} = \text{tr}(\overline{X})$  and  $\overline{\Delta} = \det(\overline{X})$ . Let  $\overline{G}_N$  denote the first fundamental solution to the  $(\overline{\tau}, \overline{\Delta})$ -recurrence (22).

Then we have the **cyclic binomial identity**

$$\begin{aligned} \overline{G}_N &= \sum_{(k_1, \dots, k_t)} \binom{N-k_2}{k_1} \binom{N-k_3}{k_2} \cdots \binom{N-k_1}{k_t} x_1^{k_1} x_2^{k_2} \cdots x_t^{k_t} a_1^{N-k_1-k_2} a_2^{N-k_2-k_3} \cdots a_t^{N-k_t-k_1} \\ &= \overline{\Delta}^{N/2} U_N \left( \frac{\overline{\tau}}{2\sqrt{\overline{\Delta}}} \right) \\ &= \sum_{k=0}^{\lfloor N/2 \rfloor} \binom{N-k}{k} \overline{\tau}^{N-2k} (-\overline{\Delta})^k, \end{aligned}$$

where  $U_N$  denotes the Chebyshev polynomial of the second kind.

*Proof.* The first equality follows by computing the symmetric trace for arbitrary  $t$  as indicated above. The second follows by induction on  $N$  using initial conditions  $\overline{G}_{-1} = 0$  and  $\overline{G}_0 = 1$ , the  $(\overline{\tau}, \overline{\Delta})$ -recurrence (22) and the Chebyshev recurrence (25). The third follows from the second by the Symmetric Trace Theorem applied to  $\overline{X} = \begin{pmatrix} 0 & 1 \\ -\overline{\Delta} & \overline{\tau} \end{pmatrix}$ , the shift matrix for the  $(\overline{\tau}, \overline{\Delta})$ -recurrence.  $\blacksquare$

Note that  $G_{-1} = 0$  and  $G_0 = 1$ , so  $G_1 = \tau$  using the  $(\tau, \Delta)$ -recurrence. This also follows directly from the condition  $k_{s-1} + k_s \leq 1$  for non-zero terms in the cyclic binomial summation above. Note also that setting all  $a_s = 1$  above gives explicit expressions for  $G_N$ .

**Example 5** Here  $N = 2$  and  $t = 3$ . Let  $A^{\text{Sym}(N)}$  denote the symmetric representation in degree  $N$  of the matrix  $A$ . From the above we have

$$\begin{aligned} G_2 &= \sum_{(k_1, k_2, k_3)} \binom{2-k_2}{k_1} \binom{2-k_3}{k_2} \binom{2-k_1}{k_3} x_1^{k_1} x_2^{k_2} x_3^{k_3} \\ &= 1 + 2x_1 + 2x_2 + 2x_3 + x_1^2 + 2x_1x_2 + 2x_1x_3 + x_2^2 + 2x_2x_3 + x_3^2 + x_1x_2x_3. \end{aligned}$$

Also  $d = 2$ , so  $\binom{N+d-1}{N} = 3$ , and  $\xi_s = \begin{pmatrix} 0 & 1 \\ x_s & 1 \end{pmatrix}$  for  $1 \leq i \leq 3$ , thus

$$X = \xi_3 \xi_2 \xi_1 = \begin{pmatrix} x_1 & x_2 + 1 \\ x_1 x_3 + x_1 & x_2 + x_3 + 1 \end{pmatrix}.$$

Now  $\xi_s^{Sym(2)} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & x_s & 1 \\ x_s^2 & 2x_s & 1 \end{pmatrix}$  for  $1 \leq s \leq 3$ , and so

$$\begin{aligned} X^{Sym(2)} &= \xi_3^{Sym(2)} \xi_2^{Sym(2)} \xi_1^{Sym(2)} \\ &= \begin{pmatrix} x_1^2 & 2x_1x_2 + 2x_1 & x_2^2 + 2x_2 + 1 \\ x_1^2x_3 + x_1^2 & x_1x_2x_3 + 2x_1x_2 + 2x_1x_3 + 2x_1 & x_2^2 + x_2x_3 + 2x_2 + x_3 + 1 \\ x_1^2x_3^2 + 2x_1^2x_3 + x_1^2 & 2x_1x_2x_3 + 2x_1x_3^2 + 2x_1x_2 + 4x_1x_3 + 2x_1 & x_3^2 + 2x_2x_3 + x_2^2 + 2x_2 + 2x_3 + 1 \end{pmatrix}. \end{aligned}$$

We check that  $G_2 = \text{tr}(X^{Sym(2)})$ , as indicated above.

We now give an expression for  $G_N$  as a quotient of two matching polynomials; this requires (29) from the next section.

**Theorem 4.5** *For  $N \geq 0$  we have*

$$G_N = \frac{\Phi_{1, Nt-2}}{\phi_{t-2}}.$$

*Proof.* Equation (29) is

$$\Phi_{i, Nt+j} = \Phi_{i,j} G_N - \Delta \phi_{i,j} G_{N-1}, \quad (26)$$

and from Example 2 we have  $\phi_{i,i-3} = 0$ . So (26) with  $j = i - 3$  gives

$$G_N = \frac{\Phi_{i, Nt+i-3}}{\Phi_{i,i-3}} = \frac{\Phi_{1, Nt-2}}{\phi_{t-2}}, \quad (27)$$

the second equality comes from putting  $i = 1$  in the first and then using (9) in the denominator.  $\blacksquare$

Finally, consider the Fibonacci sequence  $\{F_m \mid m \geq 1\} = \{1, 1, 2, 3, 5, 8, 13, 21, \dots\}$ . It is straightforward to show that the number of matchings in the path  $P_m$  with  $m-1$  edges is  $F_{m+1}$ . Now  $\Phi_{1, Nt-2}$  is the matching polynomial of the path  $P(1, Nt-2)$  which has  $(N+1)t-2$  edges and so has  $F_{(N+1)t}$  matchings. Similarly, the path whose matching polynomial is  $\phi_{t-2}$  has  $F_t$  matchings. Now, evaluating (27) above with  $N = N-1$  and  $x_s = 1$  for all  $1 \leq s \leq t$ , gives  $F_t | F_{Nt}$ , a well-known result on Fibonacci numbers, see pp. 148-9, Hardy and Wright [4]. Furthermore, we have

$$\frac{F_{(N+1)t}}{F_t} = \sum_{(k_1, \dots, k_t)} \binom{N-k_2}{k_1} \binom{N-k_3}{k_2} \cdots \binom{N-k_t}{k_{t-1}}.$$

## 5 Examples: Paths, Cycles, Trees

In this Section we express the matching polynomial of some well-known graphs in terms of the fundamental solutions to the  $(\tau, \Delta)$ -recurrence (12).

$G_N$  is the first fundamental solution to the  $(\tau, \Delta)$ -recurrence, so the initial values for  $G_N$  are

$$G_{-2} = \frac{-1}{\Delta}, \quad G_{-1} = 0, \quad G_0 = 1, \quad (\text{and } G_1 = \tau). \quad (28)$$

The second fundamental solution is  $-\Delta G_{N-1}$ .

### 5.1 Paths

$\Phi_{i, Nt+j}$  satisfies the  $(\tau, \Delta)$ -recurrence whose fundamental solutions are  $G_N$  and  $-\Delta G_{N-1}$ , thus  $\Phi_{i, Nt+j} = a G_N + b(-\Delta G_{N-1})$  for some  $a$  and  $b$ . The initial conditions for  $\Phi_{i, Nt+j}$  from (9) and for  $G_N$  from (28) give  $a = \Phi_{i,j}$  and  $b = \Phi_{i, -t+j} = \phi_{i,j}$ . Hence for  $N \geq -1$ ,

$$\Phi_{i, Nt+j} = \Phi_{i,j} G_N - \Delta \phi_{i,j} G_{N-1}. \quad (29)$$

**Example 6** Here  $i = 2$  and  $t = 3$ ,

$$\begin{aligned} N = -1 & \quad \phi_{2,2} = 1 + x_2, \\ N = 0 & \quad \phi_{2,3} = 1 + x_2 + x_3, \\ N = 0 & \quad \Phi_{2,1} = 1 + x_1 + x_2 + x_3 + x_1 x_2, \\ N = 0 & \quad \Phi_{2,2} = 1 + x_1 + 2x_2 + x_3 + x_1 x_2 + x_2^2 + x_2 x_3, \\ N = 1 & \quad \Phi_{2,3} = 1 + x_1 + 2x_2 + 2x_3 + x_1 x_2 + x_1 x_3 + x_2^2 + 2x_2 x_3 + x_3^2 + x_1 x_2 x_3. \end{aligned}$$


For  $N \geq 1$  let  $P_{Nt+j+1} = P(1, (N-1)t+j)$  be the path with  $Nt+j+1$  vertices and  $Nt+j$  edges, cyclically labelled starting with label  $x_1$ . Let  $\mathcal{P}_{Nt+j+1}(\mathbf{x}) = \mathcal{P}_{Nt+j+1} = \Phi_{1, (N-1)t+j}$  be its matching polynomial. With this notation any subscript on a  $P$ ,  $\mathcal{P}$ ,  $C$ , or  $\mathcal{C}$  refers to the number of vertices in the appropriate graph.

**Theorem 5.1** For any  $N \geq 1$  we have


$$\begin{aligned} (i) \quad \mathcal{P}_{Nt+j+1} &= \Phi_{1,j} G_{N-1} - \Delta \phi_j G_{N-2}, \\ (ii) \quad \mathcal{P}_{Nt+1} &= G_N + (\phi - \tau) G_{N-1}. \end{aligned}$$

*Proof.* The proof of (i) is clear using (29) with  $i = 1$  and  $N = N - 1$ . So (i) with  $j = 0$  gives  $\mathcal{P}_{Nt+1} = \Phi_{1,0} G_{N-1} - \Delta \phi_0 G_{N-2}$ . But  $\Phi_{1,0} = \Phi_{1,-t+t} = \phi_{1,t} = \phi_t = \phi$  and  $\phi_0 = \phi_{1,0} = 1$ , and then using the  $(\tau, \Delta)$ -recurrence for  $G_N$  gives (ii).  $\blacksquare$

**Example 7** Here  $t = 3$ ,



$$\mathcal{P}_{1.3+2+1} = 1 + 2x_1 + 2x_2 + x_3 + x_1^2 + 2x_1x_2 + x_1x_3 + x_2^2 + x_2x_3 + x_1x_2x_3.$$



$$\mathcal{P}_{2.3+1} = 1 + 2x_1 + 2x_2 + 2x_3 + x_1^2 + 2x_1x_2 + 3x_1x_3 + x_2^2 + 2x_2x_3 + x_3^2 + x_1^2x_3 + 2x_1x_2x_3 + x_1x_3^2.$$

## 5.2 Cycles

Now we identify the first and the last vertices of the path  $P(i, Nt + j)$  to form the cyclically labelled cycle  $C(i, Nt + j)$  with matching polynomial  $\Gamma_{i, Nt+j}(\mathbf{x}) = \Gamma_{i, Nt+j}$ .

By decomposing  $\Gamma_{i, Nt+j}$  at the ‘first’ edge labelled  $x_i$  we see that, *cf.* (29),

$$\begin{aligned}
\Gamma_{i, Nt+j} &= \Phi_{i+1, Nt+j} + x_i \Phi_{i+2, Nt+j-1}, \\
&= \Phi_{i+1, j} G_N - \Delta \phi_{i+1, j} G_{N-1} + x_i \{ \Phi_{i+2, j-1} G_N - \Delta \phi_{i+2, j-1} G_{N-1} \}, \\
&= \{ \Phi_{i+1, j} + x_i \Phi_{i+2, j-1} \} G_N - \Delta \{ \phi_{i+1, j} + x_i \phi_{i+2, j-1} \} G_{N-1}, \\
&= \Gamma_{i, j} G_N - \Delta \tau_{i, j} G_{N-1},
\end{aligned} \tag{30}$$

using (29) at the second line, and decomposing  $\Gamma_{i, j}$  and  $\tau_{i, j}$  at the first edge  $x_i$  at the fourth line. Also, defining  $\Gamma_{i, -t+j} = \tau_{i, j}$  ensures that (30) is true for all  $N \geq -1$ .

**Example 8** Here  $i = 2$  and  $t = 3$  again,

$$\begin{aligned}
N = -1 & \quad \tau_{2, 2} = 1, \\
N = 0 & \quad \tau_{2, 3} = 1 + x_2 + x_3, \\
N = 0 & \quad \Gamma_{2, 1} = 1 + x_1 + x_2 + x_3, \\
N = 0 & \quad \Gamma_{2, 2} = 1 + x_1 + 2x_2 + x_3 + x_1x_2 + x_2x_3, \\
N = 1 & \quad \Gamma_{2, 3} = 1 + x_1 + 2x_2 + 2x_3 + x_1x_2 + x_1x_3 + x_2^2 + x_2x_3 + x_3^2.
\end{aligned}$$

Let  $C_{Nt+j} = C(1, (N-1)t + j)$  be the cycle with  $Nt + j$  vertices and  $Nt + j$  edges in which labelling has started with  $x_1$ , and let  $\mathcal{C}_{Nt+j}(\mathbf{x}) = \mathcal{C}_{Nt+j} = \Gamma_{1, (N-1)t+j}$  be its matching polynomial. Compare with Theorem 5.1,


**Theorem 5.2** *For any  $N \geq 1$  we have*

$$\begin{aligned}
(i) \quad \mathcal{C}_{Nt+j} &= \Gamma_{1, j} G_{N-1} - \Delta \tau_j G_{N-2}, \\
(ii) \quad \mathcal{C}_{Nt} &= G_N - \Delta G_{N-2}.
\end{aligned}$$


*Proof.* The proof of (i) is clear from (30). Part (i) with  $j = 0$  gives (ii), using  $\Gamma_{1, 0} = \tau$ , and  $\tau_0 = 2$  from (8).  $\blacksquare$

**Example 9** Here  $t = 3$  again, the cycle starts at the large vertex and proceeds clockwise,





$$C_{1,3+2} = 1 + 2x_1 + 2x_2 + x_3 + x_1^2 + x_1x_2 + x_1x_3 + x_2^2 + x_2x_3$$



$$C_{2,3} = 1 + 2x_1 + 2x_2 + 2x_3 + x_1^2 + 2x_1x_2 + 2x_1x_3 + x_2^2 + 2x_2x_3 + x_3^2 + 2x_1x_2x_3.$$

For a fixed  $t \geq 1$  write  $\widehat{\mathcal{P}}_N = \mathcal{P}_{Nt+1}$  and  $\widehat{\mathcal{C}}_N = \mathcal{C}_{Nt}$ . We now express  $G_N$ ,  $\widehat{\mathcal{P}}_N$ , and  $\widehat{\mathcal{C}}_N$  in terms of Chebyshev polynomials.

It is well-known that, in one variable  $x$ , the matching polynomial of the path  $P_{2m}$  is related to  $U_{2m}$  as follows

$$\mathcal{M}(P_{2m}, x) = (-1)^m x^m U_{2m} \left( \frac{i}{2\sqrt{x}} \right),$$

and, for  $P_{2m-1}$  we have

$$\mathcal{M}(P_{2m-1}, x) = (-1)^m x^m \left[ U_{2m} \left( \frac{i}{2\sqrt{x}} \right) + U_{2m-2} \left( \frac{i}{2\sqrt{x}} \right) \right],$$

where  $i = \sqrt{-1}$ . Also, for the matching polynomials  $\mathcal{M}(C_{2m})$  and  $\mathcal{M}(C_{2m-1})$  of the cycles  $C_{2m}$  and  $C_{2m-1}$  there are similar formulas but with a factor of 2 on the right-hand side where  $U$  is replaced by  $T$ . See Theorem 3 of Godsil and Gutman [3], and Theorems 9 and 11 of Farrell [1].

Now Theorem 4.4 modified for  $G_N$  gives

$$G_N = \Delta^{N/2} U_N \left( \frac{\tau}{2\sqrt{\Delta}} \right). \quad (31)$$

Formulas for  $\widehat{\mathcal{P}}_N$  and  $\widehat{\mathcal{C}}_N$  in terms of  $U_N$  and  $T_N$  are given below, where the variable  $t$  is suppressed.

**Theorem 5.3** *For any  $N \geq 1$  we have*

$$(i) \quad \widehat{\mathcal{P}}_N = \Delta^{N/2} \left\{ U_N \left( \frac{\tau}{2\sqrt{\Delta}} \right) + \left( \frac{\phi - \tau}{\sqrt{\Delta}} \right) U_{N-1} \left( \frac{\tau}{2\sqrt{\Delta}} \right) \right\},$$

$$(ii) \quad \widehat{\mathcal{C}}_N = 2\Delta^{N/2} T_N \left( \frac{\tau}{2\sqrt{\Delta}} \right).$$

*Proof.* (i) This follows from Theorem 5.1(ii) and (31).

(ii) From Theorem 5.2(ii) we have  $\widehat{\mathcal{C}}_N = G_N - \Delta G_{N-2}$ , and now the well-known relation  $2T_N = U_N - U_{N-2}$  between the two types of Chebyshev polynomials and (31) gives the result.  $\blacksquare$

Expressions for  $G_N$ ,  $\widehat{\mathcal{P}}_N$ , and  $\widehat{\mathcal{C}}_N$  for  $N = 0, 1, 2, 3$ , and 4 are given below

$$\begin{array}{lll} G_0 = 1 & \widehat{\mathcal{P}}_0 = 1 & \widehat{\mathcal{C}}_0 = 2 \\ G_1 = \tau & \widehat{\mathcal{P}}_1 = \phi & \widehat{\mathcal{C}}_1 = \tau \\ G_2 = \tau^2 - \Delta & \widehat{\mathcal{P}}_2 = \phi\tau - \Delta & \widehat{\mathcal{C}}_2 = \tau^2 - 2\Delta \\ G_3 = \tau^3 - 2\tau\Delta & \widehat{\mathcal{P}}_3 = \phi\tau^2 - \phi\Delta - \tau\Delta & \widehat{\mathcal{C}}_3 = \tau^3 - 3\tau\Delta \\ G_4 = \tau^4 - 3\tau^2\Delta + \Delta^2 & \widehat{\mathcal{P}}_4 = \phi\tau^3 - 2\phi\tau\Delta - \tau^2\Delta + \Delta^2 & \widehat{\mathcal{C}}_4 = \tau^4 - 4\tau^2\Delta + 2\Delta^2. \end{array}$$

### 5.3 Trees

Here we consider cyclically labelled trees.

First let us extend the definition of a cyclically labelled path to include the path of Fig. 1, and the graph  $P_1$  with one vertex and no edges.

A tree is a connected simple graph with no cycles, and a rooted tree is a tree in which some vertex of degree 1 has been specified to be the root,  $r$ . Given any rooted tree, let us label its edges by first labelling the edge incident to  $r$  with  $x_i$ . Then label all edges incident to this edge with  $x_{i+1}$ , then label all edges incident to these edges with  $x_{i+2}$ , and so on until label  $x_t$  has been used. Then label with the ordered set  $\{x_1, \dots, x_t\}$  in a similar

manner to before, repeating cyclically until all edges have been labelled,..., and so on. Let  $T$  denote such a cyclically labelled tree, see Fig. 5 for an example with  $i = 2$  and  $t = 3$ .




Fig. 5: A cyclically labelled tree with  $i = 2$  and  $t = 3$ .

We may draw any such  $T$  with  $r$  as the leftmost vertex. Then we place the other vertices of  $T$  from ‘left to right’ according to their distance from  $r$ , *i.e.*, if a vertex  $v_1$  is at distance  $d_1$  from  $r$  and vertex  $v_2$  is at distance  $d_2$  from  $r$  where  $d_2 > d_1$ , then  $v_2$  is placed to the right of  $v_1$ .

Paths in  $T$  are of two types: (I) A path that always moves from left to right (a path that always moves from right to left can be thought of one that always moves from left to right): such a path is clearly cyclically labelled; or (II) a path that moves first from right to left and then from left to right; such a path must pass through at least one vertex of degree  $\geq 3$ , *i.e.*, a vertex where  $T$  ‘branches’.

Let  $V$  denote the set of vertices of degree  $\geq 3$  in  $T$ , and let  $v \in V$  be arbitrary of degree  $\deg(v)$ . Vertex  $v$  has 1 edge to its left and  $\deg(v) - 1 \geq 2$  edges to its right. Let  $H_v$  be the subgraph of  $T$  that consists of the ‘last’  $\deg(v) - 2 \geq 1$  edges as we rotate clockwise around  $v$ . Thus  $H_v$  is the star  $K_{1, \deg(v)-2}$  centered at  $v$ . Set  $H = \cup_{v \in V} H_v$ .

**Lemma 5.4** *The forest  $T - H$  is a union of cyclically labelled paths.*

*Proof.* We show that  $T - H$  does not contain a path of type (II). Suppose it does contain a path of type (II), then this path must pass through some

vertex  $v \in V$ . So 2 edges incident to  $v$  and to the right of  $v$  lie in this path and so lie in  $T - H$ , a contradiction because  $T - H$  contains only 1 edge incident to  $v$  and to the right of  $v$ . Thus  $T - H$  is a union of paths of type (I), each of which is a cyclically labelled path. ■

Thus  $T - H$  is a union of cyclically labelled paths, and so  $T - H - \overline{M}_H$  is also, for every matching  $M_H$  of  $H$ . We know the matching polynomial of any cyclically labelled path, so we can decompose the matching polynomial of  $T$ ,  $\mathcal{M}(T, \mathbf{x})$ , at  $H$ , according to Theorem 1.1,

$$\mathcal{M}(T, \mathbf{x}) = \sum_{M_H} M_H(\mathbf{x}) \mathcal{M}(T - H - \overline{M}_H, \mathbf{x}),$$

where the summation is over every matching  $M_H$  of  $H$ .

**Example 10** See Fig. 5.



$H$  has 6 matchings with weights:  $1, x_1, x_1, x_3, x_1x_3$ , and  $x_1x_3$ . Thus there are 6 terms in the decomposition, and  $\mathcal{M}(T, \mathbf{x})$  is the sum of the following 6 terms:

$$\begin{aligned} & 1 \cdot \phi_1 \phi_{2,3} \Phi_{2,1} + x_1 \cdot \phi_1 \phi_{2,2} \phi_{2,3} + x_1 \cdot \phi_1 \phi_{2,2} \phi_{3,3} + x_3 \cdot \phi_1 \phi_{2,3} + x_1 x_3 \cdot \phi_{2,3} + x_1 x_3 \cdot \phi_{3,3} \\ & = 1 + 4x_1 + 2x_2 + 3x_3 + 3x_1^2 + 7x_1x_2 + 8x_1x_3 + x_2^2 + 3x_2x_3 + 2x_3^2 \\ & \quad + 5x_1^2x_2 + 3x_1^2x_3 + 3x_1x_2^2 + 7x_1x_2x_3 + 4x_1x_3^2 + 2x_1^2x_2^2 + 3x_1^2x_2x_3. \end{aligned}$$

**Acknowledgement:** We thank the referees for some helpful comments.

## References

- [1] E.J. Farrell, *An Introduction to Matching Polynomials*, Jour. Comb. Theory, Series B, **27**, pp. 75–86, 1979.
- [2] W. Fulton, J. Harris, *Representation Theory, A First Course*, Graduate Texts in Mathematics, **129**, Springer-Verlag, 1991.

- [3] C.D. Godsil, I. Gutman, *On the Theory of the Matching Polynomial*, Jour. Graph. Theory, **5**, pp. 137–144, 1981.
- [4] G.H. Hardy, E.M. Wright, *An Introduction to the Theory of Numbers*, 5th Edition, Oxford University Press, 1979.
- [5] T.A. Springer, *Invariant Theory*, Lecture Notes in Mathematics, **585**, Springer-Verlag, 1977.